

Disclaimer

The following report(s) provides findings from an FDA-initiated query using Sentinel. While Sentinel queries may be undertaken to assess potential medical product safety risks, they may also be initiated for various other reasons. Some examples include determining a rate or count of an identified health outcome of interest, examining medical product use, exploring the feasibility of future, more detailed analyses within Sentinel, and seeking to better understand Sentinel capabilities.

Data obtained through Sentinel are intended to complement other types of evidence such as preclinical studies, clinical trials, postmarket studies, and adverse event reports, all of which are used by FDA to inform regulatory decisions regarding medical product safety. The information contained in this report is provided as part of FDA's commitment to place knowledge acquired from Sentinel in the public domain as soon as possible. Any public health actions taken by FDA regarding products involved in Sentinel queries will continue to be communicated through existing channels.

FDA wants to emphasize that the fact that FDA has initiated a query involving a medical product and is reporting findings related to that query does not mean that FDA is suggesting health care practitioners should change their prescribing practices for the medical product or that patients taking the medical product should stop using it. Patients who have questions about the use of an identified medical product should contact their health care practitioners.

The following report contains a description of the request, request specifications, and results from the modular program run(s).

If you are using a web page screen reader and are unable to access this document, please contact the Sentinel Operations Center for assistance at info@sentinelsystem.org.



Overview

Date Run: October 25, 2017

Request Description: The purpose of this report was to compare the frequency of diagnoses for intracranial hemorrhage using International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM) versus International Classification of Diseases, Tenth Revision, Clinical Modification (ICD-10-CM) diagnosis codes. ICD-10-CM code definitions were determined by mapping from ICD-9-CM code definitions using the Centers for Medicare and Medicaid Services (CMS) General Equivalence Mappings (GEMs). Forward-backward mapping (FBM) was used to map ICD-9-CM to ICD-10-CM codes.¹

Sentinel Modular Program Tool Used: Cohort Identification and Descriptive Analysis (CIDA) tool, version 5.0.5

<u>Data Source:</u> This request was run against data from 12 Data Partners contributing to the Sentinel Distributed Database (SDD). Data from October 1, 2010 to September 30, 2016 were included in this report. The report includes three separate time periods: 1) October 1, 2010 to September 30, 2016; 2) April 1, 2015 to September 30, 2015; and 3) April 1, 2016 to September 30, 2016. This request was distributed to Data Partners on October 25, 2017. See Appendix A for a list of dates of available data for each Data Partner.

<u>Study Design:</u> We examined the incidence and prevalence across the ICD-9-CM era (October 2010 - September 2015) and ICD-10-CM era (October 2015 - September 2016) in the United States. Incidence was additionally evaluated from April 2015 to September 2015 and April 2016 to September 2016. See Appendix B for specific codes used to define intracranial hemorrhage in this request.

<u>Cohort Eligibility Criteria:</u> Members included in the cohort were required to be continuously enrolled in health plans with medical and drug coverage for at least six months (183 days) before their diagnosis date, during which gaps in coverage of up to 45 days were allowed. The following age groups were included in the cohort: 21-64, 65-74, 75-84, and 85-99 years.

<u>Incident Cohorts:</u> Members included in the incident cohorts were required to be continuously enrolled in health plans with medical and drug coverage for at least 183 days prior to intracranial hemorrhage diagnosis, during which gaps in coverage of up to 45 days were allowed. Incident intracranial hemorrhage was defined as no previous intracranial hemorrhage diagnosis in the 183 days preceding the index date with respect to ICD-9-CM and ICD-10-CM codes.

<u>Prevalent Cohorts:</u> There was no enrollment time requirement for members in the prevalent cohorts. All qualifying diagnosis codes that occurred between October 1, 2010 and September 30, 2016 were included.

Please see Appendix C for detailed specifications of parameters used in the analyses for this request.

<u>Limitations:</u> Algorithms used to define outcomes are imperfect; thus, it is possible that there may be misclassification. Therefore, data should be interpreted with this limitation in mind.

Notes: Please contact the Sentinel Operations Center (info@sentinelsystem.org) for questions and to provide comments/suggestions for future enhancements to this document.

¹Fung, K. W., et al. (2016). "Preparing for the ICD-10-CM Transition: Automated Methods for Translating ICD Codes in Clinical Phenotype Definitions." EGEMS (Wash DC) 4(1): 1211.



	Table of Contents
Glossary	List of Terms Found in this Report and their Definitions
Table 1	Comparison of Incident Intracranial Hemorrhage Diagnoses in the International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM) and International Classification of Diseases, Tenth Revision, Clinical Modification (ICD-10-CM) Eras (April 1, 2015 - September 30, 2015 and April 1, 2016 - September 30, 2016)
Figure 1	Incidence of Intracranial Hemorrhage Diagnoses, per 1,000 Eligible Members from October 2010 - September 2016 by Code Type, 183-Day Washout
Figure 2	Prevalence of Intracranial Hemorrhage Diagnoses per 1,000 Eligible Members from October 2010 - September 2016 by Code Type, 0-Day Washout
Appendix A	Dates of Available Data for Each Data Partner as of Request Distribution Date (October 25, 2017)
Appendix B	List of International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM) and International Classification of Diseases, Tenth Revision, Clinical Modification (ICD-10-CM) Codes Used to Define Intracranial Hemorrhage
Appendix C	Specifications for Parameters for this Request



Glossary of Terms for Analyses Using Cohort Identification and Descriptive Analysis (CIDA) Tool*

Amount Supplied - number of units (pills, tablets, vials) dispensed. Net amount per NDC per dispensing.

Blackout Period - number of days at the beginning of a treatment episode that events are to be ignored. If an event occurs during the blackout period, the episode is excluded.

Care Setting - type of medical encounter or facility where the exposure, event, or condition code was recorded. Possible care settings include: Inpatient Hospital Stay (IP), Non-Acute Institutional Stay (IS), Emergency Department (ED), Ambulatory Visit (AV), and Other Ambulatory Visit (OA). For laboratory results, possible care settings include: Emergency Department (E), Home (H), Inpatient (I), Outpatient (O), or Unknown or Missing (U). The Care Setting, along with the Principal Diagnosis Indicator, forms the Care Setting/PDX parameter.

Ambulatory Visit (AV) - includes visits at outpatient clinics, same-day surgeries, urgent care visits, and other same-day ambulatory hospital encounters, but excludes emergency department encounters.

Emergency Department (ED) - includes ED encounters that become inpatient stays (in which case inpatient stays would be a separate encounter). Excludes urgent care visits.

Inpatient Hospital Stay (IP) - includes all inpatient stays, same-day hospital discharges, hospital transfers, and acute hospital care where the discharge is after the admission date.

Non-Acute Institutional Stay (IS) - includes hospice, skilled nursing facility (SNF), rehab center, nursing home, residential, overnight non-hospital dialysis and other non-hospital stays.

Other Ambulatory Visit (OA) - includes other non overnight AV encounters such as hospice visits, home health visits, skilled nursing facility visits, other non-hospital visits, as well as telemedicine, telephone and email consultations.

Cohort Definition (drug/exposure) - indicates how the cohort will be defined: 01: Cohort includes only the first valid treatment episode during the query period; 02: Cohort includes all valid treatment episodes during the query period; 03: Cohort includes all valid treatment episodes during the query period until an event occurs.

Days Supplied - number of days supplied for all dispensings in qualifying treatment episodes.

Eligible Members - number of members eligible for an incident treatment episode (defined by the drug/exposure and event washout periods) with drug and medical coverage during the query period.

Enrollment Gap - number of days allowed between two consecutive enrollment periods without breaking a "continuously enrolled" sequence.

Episodes - treatment episodes; length of episode is determined by days supplied in one dispensing or consecutive dispensings bridged by the episode gap.

Episode Gap - number of days allowed between two (or more) consecutive exposures (dispensings/procedures) to be considered the same treatment episode.

Event Deduplication - specifies how events are counted by the MP algorithm: 0: Counts all occurrences of an HOI during an exposure episode; 1: de-duplicates occurrences of the same HOI code and code type on the same day; 2: de-duplicates occurrences of the same HOI group on the same day (e.g., de-duplicates at the group level).

Exposure Episode Length - number of days after exposure initiation that is considered "exposed time."

Exposure Extension Period - number of days post treatment period in which the outcomes/events are counted for a treatment episode. Extensions days are added after any episode gaps have been bridged

Lookback Period - number of days wherein a member is required to have evidence of pre-existing condition (diagnosis/procedure/drug dispensing).



Maximum Episode Duration - truncates exposure episodes after a requester-specified number of exposed days. Applied after any gaps are bridged and extension days added to the length of the exposure episode.

Member-Years - sum of all days of enrollment with medical and drug coverage in the query period preceded by an exposure washout period all divided by 365.25.

Minimum Days Supplied - specifies a minimum number of days in length of the days supplied for the episode to be considered.

Minimum Episode Duration - specifies a minimum number of days in length of the episode for it to be considered. Applied after any gaps are bridged and extension days added to the length of the exposure episode.

Monitoring Period - used to define time periods of interest for both sequential analysis and simple cohort characterization requests.

Principal Diagnosis (PDX) - diagnosis or condition established to be chiefly responsible for admission of the patient to the hospital. 'P' = principal diagnosis, 'S' = secondary diagnosis, 'X' = unspecified diagnosis, '.' = blank. Along with the Care Setting values, forms the Caresetting/PDX parameter.

Query Period - period in which the modular program looks for exposures and outcomes of interest.

Treatment Episode Truncation Indicator - indicates whether the exposure episode will be truncated at the occurrence of a requester-specified code.

Washout Period (drug/exposure) - number of days a user is required to have no evidence of prior exposure (drug dispensing/procedure) and continuous drug and medical coverage prior to an incident treatment episode.

Washout Period (event/outcome) - number of days a user is required to have no evidence of a prior event (procedure/diagnosis) and continuous drug and medical coverage prior to an incident treatment episode.

Years at Risk - number of days supplied plus any episode gaps and exposure extension periods all divided by 365.25.

*all terms may not be used in this report



Table 1. Comparison of Incident* Intracranial Hemorrhage Diagnoses in the International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM) and International Classification of Diseases, Tenth Revision, Clinical Modification (ICD-10-CM) Eras (April 1, 2015 - September 30, 2015 and April 1, 2016 - September 30, 2016)

	Members with Diagnosis	Eligible Members	Members with Diagnosis per 1,000 Eligible Members
Intracranial Hemorrhage			
ICD-9-CM: April 1, 2015 - September 30, 2015			
	27,882	32,383,988	0.86
ICD-10-CM: April 1, 2016 - September 30, 2016			
	28,122	33,760,393	0.83

^{*}Incidence defined by a 183 day washout



Figure 1. Incidence of Intracranial Hemorrhage Diagnoses per 1,000 Eligible Members from October 2010 - September 2016 by Code Type, 183-Day Washout







Figure 2. Prevalence of Intracranial Hemorrhage Diagnoses per 1,000 Eligible Members from October 2010 - September 2016 by Code Type, 0-Day Washout





Appendix A. Dates of Available Data for Each Data Partner as of Request Distribution Date (October 25, 2017)

DP ID	Start Date ¹	End Date ¹
DP0001	10/1/2010	6/30/2016
DP0002	10/1/2010	6/30/2016
DP0003	10/1/2010	9/30/2016
DP0004	10/1/2010	9/30/2016
DP0005	10/1/2010	9/30/2016
DP0006	10/1/2010	9/30/2016
DP0007	10/1/2010	9/30/2016
DP0008	10/1/2010	9/30/2016
DP0009	10/1/2010	9/30/2016
DP0010	10/1/2010	9/30/2016
DP0011	10/1/2010	9/30/2016
DP0012	10/1/2010	9/30/2016

¹The start and end dates are based on the minimum and maximum dates within each DP. The month with the maximum date must have at least 80% of the number of records in the previous month.



Code	Description	Code Type
	ICD-9-CM	
430	Subarachnoid hemorrhage	ICD-9-CM
431	Intracerebral hemorrhage	ICD-9-CM
432	Other and unspecified intracranial hemorrhage	ICD-9-CM
432.0	Nontraumatic extradural hemorrhage	ICD-9-CM
432.1	Subdural hemorrhage	ICD-9-CM
432.9	Unspecified intracranial hemorrhage	ICD-9-CM
852.0	Subarachnoid hemorrhage following injury without mention of open intracranial wound	ICD-9-CM
852.00	Subarachnoid hemorrhage following injury, without mention of open intracranial wound, unspecified state of consciousness	ICD-9-CM
852.01	Subarachnoid hemorrhage following injury, without mention of open intracranial wound, no loss of consciousness	ICD-9-CM
852.02	Subarachnoid hemorrhage following injury, without mention of open intracranial wound, brief (less than 1 hour) loss of consciousness	ICD-9-CM
852.03	Subarachnoid hemorrhage following injury, without mention of open intracranial wound, moderate (1-24 hours) loss of consciousness	ICD-9-CM
852.04	Subarachnoid hemorrhage following injury, without mention of open intracranial wound, prolonged (more than 24 hours) loss of consciousness and return to preexisting conscious level	ICD-9-CM
852.05	Subarachnoid hemorrhage following injury, without mention of open intracranial wound, prolonged (more than 24 hours) loss of consciousness, without return to preexisting conscious level	ICD-9-CM
852.06	Subarachnoid hemorrhage following injury, without mention of open intracranial wound, loss of consciousness of unspecified duration	ICD-9-CM
852.09	Subarachnoid hemorrhage following injury, without mention of open intracranial wound, unspecified concussion	ICD-9-CM
852.2	Subdural hemorrhage following injury without mention of open intracranial wound	ICD-9-CM
852.20	Subdural hemorrhage following injury, without mention of open intracranial wound, unspecified state of consciousness	ICD-9-CM
852.21	Subdural hemorrhage following injury, without mention of open intracranial wound, no loss of consciousness	ICD-9-CM
852.22	Subdural hemorrhage following injury, without mention of open intracranial wound, brief (less than one hour) loss of consciousness	ICD-9-CM
852.23	Subdural hemorrhage following injury, without mention of open intracranial wound, moderate (1-24 hours) loss of consciousness	ICD-9-CM
852.24	Subdural hemorrhage following injury, without mention of open intracranial wound, prolonged (more than 24 hours) loss of consciousness and return to pre-existing conscious level	ICD-9-CM
852.25	Subdural hemorrhage following injury, without mention of open intracranial wound, prolonged (more than 24 hours) loss of consciousness, without return to pre-existing conscious level	ICD-9-CM
852.26	Subdural hemorrhage following injury, without mention of open intracranial wound, loss of consciousness of unspecified duration	ICD-9-CM



Code	Description	Code Type
52.29	Subdural hemorrhage following injury, without mention of open intracranial wound, unspecified concussion	ICD-9-CM
52.4	Extradural hemorrhage following injury without mention of open intracranial wound	ICD-9-CM
52.40	Extradural hemorrhage following injury, without mention of open intracranial wound, unspecified state of consciousness	ICD-9-CM
52.41	Extradural hemorrhage following injury, without mention of open intracranial wound, no loss of consciousness	ICD-9-CM
52.42	Extradural hemorrhage following injury, without mention of open intracranial wound, brief (less than 1 hour) loss of consciousness	ICD-9-CM
52.43	Extradural hemorrhage following injury, without mention of open intracranial wound, moderate (1-24 hours) loss of consciousness	ICD-9-CM
52.44	Extradural hemorrhage following injury, without mention of open intracranial wound, prolonged (more than 24 hours) loss of consciousness and return to pre-existing conscious level	ICD-9-CM
52.45	Extradural hemorrhage following injury, without mention of open intracranial wound, prolonged (more than 24 hours) loss of consciousness, without return to pre-existing conscious level	ICD-9-CM
52.46	Extradural hemorrhage following injury, without mention of open intracranial wound, loss of consciousness of unspecified duration	ICD-9-CM
52.49	Extradural hemorrhage following injury, without mention of open intracranial wound, unspecified concussion	ICD-9-CM
53.0	Other and unspecified intracranial hemorrhage following injury, without mention of open intracranial wound	ICD-9-CM
53.00	Other and unspecified intracranial hemorrhage following injury, without mention of open intracranial wound, unspecified state of consciousness	ICD-9-CM
53.01	Other and unspecified intracranial hemorrhage following injury, without mention of open intracranial wound, no loss of consciousness	ICD-9-CM
53.02	Other and unspecified intracranial hemorrhage following injury, without mention of open intracranial wound, brief (less than 1 hour) loss of consciousness	ICD-9-CM
53.03	Other and unspecified intracranial hemorrhage following injury, without mention of open intracranial wound, moderate (1-24 hours) loss of consciousness	ICD-9-CM
53.04	Other and unspecified intracranial hemorrhage following injury, without mention of open intracranial wound, prolonged (more than 24 hours) loss of consciousness and return to preexisting conscious level	ICD-9-CM
53.05	Other and unspecified intracranial hemorrhage following injury. Without mention of open intracranial wound, prolonged (more than 24 hours) loss of consciousness, without return to pre-existing conscious level	ICD-9-CM
53.06	Other and unspecified intracranial hemorrhage following injury, without mention of open intracranial wound, loss of consciousness of unspecified duration	ICD-9-CM
53.09	Other and unspecified intracranial hemorrhage following injury, without mention of open intracranial wound, unspecified concussion	ICD-9-CM
	ICD-10-CM	
0.00	Nontraumatic subarachnoid hemorrhage from unspecified carotid siphon and	ICD-10-CM

bifurcation



Code	Description	Code Type
160.01	Nontraumatic subarachnoid hemorrhage from right carotid siphon and bifurcation	ICD-10-CM
160.02	Nontraumatic subarachnoid hemorrhage from left carotid siphon and bifurcation	ICD-10-CM
160.10	Nontraumatic subarachnoid hemorrhage from unspecified middle cerebral artery	ICD-10-CM
160.11	Nontraumatic subarachnoid hemorrhage from right middle cerebral artery	ICD-10-CM
160.12	Nontraumatic subarachnoid hemorrhage from left middle cerebral artery	ICD-10-CM
160.2	Nontraumatic subarachnoid hemorrhage from anterior communicating artery	ICD-10-CM
160.30	Nontraumatic subarachnoid hemorrhage from unspecified posterior communicating artery	ICD-10-CM
160.31	Nontraumatic subarachnoid hemorrhage from right posterior communicating artery	ICD-10-CM
160.32	Nontraumatic subarachnoid hemorrhage from left posterior communicating artery	ICD-10-CM
160.4	Nontraumatic subarachnoid hemorrhage from basilar artery	ICD-10-CM
160.50	Nontraumatic subarachnoid hemorrhage from unspecified vertebral artery	ICD-10-CM
160.51	Nontraumatic subarachnoid hemorrhage from right vertebral artery	ICD-10-CM
160.52	Nontraumatic subarachnoid hemorrhage from left vertebral artery	ICD-10-CM
160.6	Nontraumatic subarachnoid hemorrhage from other intracranial arteries	ICD-10-CM
160.7	Nontraumatic subarachnoid hemorrhage from unspecified intracranial artery	ICD-10-CM
160.8	Other nontraumatic subarachnoid hemorrhage	ICD-10-CM
160.9	Nontraumatic subarachnoid hemorrhage, unspecified	ICD-10-CM
161.0	Nontraumatic intracerebral hemorrhage in hemisphere, subcortical	ICD-10-CM
161.1	Nontraumatic intracerebral hemorrhage in hemisphere, cortical	ICD-10-CM
161.2	Nontraumatic intracerebral hemorrhage in hemisphere, unspecified	ICD-10-CM
161.3	Nontraumatic intracerebral hemorrhage in brain stem	ICD-10-CM
161.4	Nontraumatic intracerebral hemorrhage in cerebellum	ICD-10-CM
161.5	Nontraumatic intracerebral hemorrhage, intraventricular	ICD-10-CM
161.6	Nontraumatic intracerebral hemorrhage, multiple localized	ICD-10-CM
161.8	Other nontraumatic intracerebral hemorrhage	ICD-10-CM
161.9	Nontraumatic intracerebral hemorrhage, unspecified	ICD-10-CM
162.00	Nontraumatic subdural hemorrhage, unspecified	ICD-10-CM
162.01	Nontraumatic acute subdural hemorrhage	ICD-10-CM
162.02	Nontraumatic subacute subdural hemorrhage	ICD-10-CM
162.03	Nontraumatic chronic subdural hemorrhage	ICD-10-CM
162.1	Nontraumatic extradural hemorrhage	ICD-10-CM
162.9	Nontraumatic intracranial hemorrhage, unspecified	ICD-10-CM
S06.340A	Traumatic hemorrhage of right cerebrum without loss of consciousness, initial encounter	ICD-10-CM
S06.341A	Traumatic hemorrhage of right cerebrum with loss of consciousness of 30 minutes or less, initial encounter	ICD-10-CM
S06.342A	Traumatic hemorrhage of right cerebrum with loss of consciousness of 31 minutes to 59 minutes, initial encounter	ICD-10-CM
S06.343A	Traumatic hemorrhage of right cerebrum with loss of consciousness of 1 hours to 5 hours 59 minutes, initial encounter	ICD-10-CM
S06.344A	Traumatic hemorrhage of right cerebrum with loss of consciousness of 6 hours to 24 hours, initial encounter	ICD-10-CM



Code	Description	Code Type
S06.345A	Traumatic hemorrhage of right cerebrum with loss of consciousness greater than 24 hours with return to pre-existing conscious level, initial encounter	ICD-10-CM
S06.346A	Traumatic hemorrhage of right cerebrum with loss of consciousness greater than 24 hours without return to pre-existing conscious level with patient surviving, initial encounter	ICD-10-CM
S06.347A	Traumatic hemorrhage of right cerebrum with loss of consciousness of any duration with death due to brain injury prior to regaining consciousness, initial encounter	ICD-10-CM
S06.348A	Traumatic hemorrhage of right cerebrum with loss of consciousness of any duration with death due to other cause prior to regaining consciousness, initial encounter	ICD-10-CM
S06.349A	Traumatic hemorrhage of right cerebrum with loss of consciousness of unspecified duration, initial encounter	ICD-10-CM
S06.350A	Traumatic hemorrhage of left cerebrum without loss of consciousness, initial encounter	ICD-10-CM
S06.351A	Traumatic hemorrhage of left cerebrum with loss of consciousness of 30 minutes or less, initial encounter	ICD-10-CM
S06.352A	Traumatic hemorrhage of left cerebrum with loss of consciousness of 31 minutes to 59 minutes, initial encounter	ICD-10-CM
S06.353A	Traumatic hemorrhage of left cerebrum with loss of consciousness of 1 hours to 5 hours 59 minutes, initial encounter	ICD-10-CM
S06.354A	Traumatic hemorrhage of left cerebrum with loss of consciousness of 6 hours to 24 hours, initial encounter	ICD-10-CM
S06.355A	Traumatic hemorrhage of left cerebrum with loss of consciousness greater than 24 hours with return to pre-existing conscious level, initial encounter	ICD-10-CM
S06.356A	Traumatic hemorrhage of left cerebrum with loss of consciousness greater than 24 hours without return to pre-existing conscious level with patient surviving, initial encounter	ICD-10-CM
S06.357A	Traumatic hemorrhage of left cerebrum with loss of consciousness of any duration with death due to brain injury prior to regaining consciousness, initial encounter	ICD-10-CM
S06.358A	Traumatic hemorrhage of left cerebrum with loss of consciousness of any duration with death due to other cause prior to regaining consciousness, initial encounter	ICD-10-CM
S06.359A	Traumatic hemorrhage of left cerebrum with loss of consciousness of unspecified duration, initial encounter	ICD-10-CM
S06.360A	Traumatic hemorrhage of cerebrum, unspecified, without loss of consciousness, initial encounter	ICD-10-CM
S06.361A	Traumatic hemorrhage of cerebrum, unspecified, with loss of consciousness of 30 minutes or less, initial encounter	ICD-10-CM
S06.362A	Traumatic hemorrhage of cerebrum, unspecified, with loss of consciousness of 31 minutes to 59 minutes, initial encounter	ICD-10-CM
S06.363A	Traumatic hemorrhage of cerebrum, unspecified, with loss of consciousness of 1 hours to 5 hours 59 minutes, initial encounter $\frac{1}{2}$	ICD-10-CM
S06.364A	Traumatic hemorrhage of cerebrum, unspecified, with loss of consciousness of 6 hours to 24 hours, initial encounter	ICD-10-CM
S06.365A	Traumatic hemorrhage of cerebrum, unspecified, with loss of consciousness greater than 24 hours with return to pre-existing conscious level, initial encounter	ICD-10-CM



Code	Description	Code Type
S06.366A	Traumatic hemorrhage of cerebrum, unspecified, with loss of consciousness greater than 24 hours without return to pre-existing conscious level with patient surviving, initial encounter	ICD-10-CM
S06.367A	Traumatic hemorrhage of cerebrum, unspecified, with loss of consciousness of any duration with death due to brain injury prior to regaining consciousness, initial encounter	ICD-10-CM
S06.368A	Traumatic hemorrhage of cerebrum, unspecified, with loss of consciousness of any duration with death due to other cause prior to regaining consciousness, initial encounter	ICD-10-CM
S06.369A	Traumatic hemorrhage of cerebrum, unspecified, with loss of consciousness of unspecified duration, initial encounter	ICD-10-CM
S06.4X0A	Epidural hemorrhage without loss of consciousness, initial encounter	ICD-10-CM
S06.4X1A	Epidural hemorrhage with loss of consciousness of 30 minutes or less, initial encounter	ICD-10-CM
S06.4X2A	Epidural hemorrhage with loss of consciousness of 31 minutes to 59 minutes, initial encounter	ICD-10-CM
S06.4X3A	Epidural hemorrhage with loss of consciousness of 1 hour to 5 hours 59 minutes, initial encounter	ICD-10-CM
S06.4X4A	Epidural hemorrhage with loss of consciousness of 6 hours to 24 hours, initial encounter	ICD-10-CM
S06.4X5A	Epidural hemorrhage with loss of consciousness greater than 24 hours with return to pre-existing conscious level, initial encounter	ICD-10-CM
S06.4X6A	Epidural hemorrhage with loss of consciousness greater than 24 hours without return to pre-existing conscious level with patient surviving, initial encounter	ICD-10-CM
S06.4X7A	Epidural hemorrhage with loss of consciousness of any duration with death due to brain injury prior to regaining consciousness, initial encounter	ICD-10-CM
S06.4X8A	Epidural hemorrhage with loss of consciousness of any duration with death due to other causes prior to regaining consciousness, initial encounter	ICD-10-CM
S06.4X9A	Epidural hemorrhage with loss of consciousness of unspecified duration, initial encounter	ICD-10-CM
S06.5X0A	Traumatic subdural hemorrhage without loss of consciousness, initial encounter	ICD-10-CM
S06.5X1A	Traumatic subdural hemorrhage with loss of consciousness of 30 minutes or less, initial encounter	ICD-10-CM
S06.5X2A	Traumatic subdural hemorrhage with loss of consciousness of 31 minutes to 59 minutes, initial encounter	ICD-10-CM
S06.5X3A	Traumatic subdural hemorrhage with loss of consciousness of 1 hour to 5 hours 59 minutes, initial encounter	ICD-10-CM
S06.5X4A	Traumatic subdural hemorrhage with loss of consciousness of 6 hours to 24 hours, initial encounter	ICD-10-CM
S06.5X5A	Traumatic subdural hemorrhage with loss of consciousness greater than 24 hours with return to pre-existing conscious level, initial encounter	ICD-10-CM
S06.5X6A	Traumatic subdural hemorrhage with loss of consciousness greater than 24 hours without return to pre-existing conscious level with patient surviving, initial encounter	ICD-10-CM
S06.5X7A	Traumatic subdural hemorrhage with loss of consciousness of any duration with death due to brain injury before regaining consciousness, initial encounter	ICD-10-CM



Code	Description	Code Type
S06.5X8A	Traumatic subdural hemorrhage with loss of consciousness of any duration with death due to other cause before regaining consciousness, initial encounter	ICD-10-CM
S06.5X9A	Traumatic subdural hemorrhage with loss of consciousness of unspecified duration, initial encounter	ICD-10-CM
S06.6X0A	Traumatic subarachnoid hemorrhage without loss of consciousness, initial encounter	ICD-10-CM
S06.6X1A	Traumatic subarachnoid hemorrhage with loss of consciousness of 30 minutes or less, initial encounter	ICD-10-CM
S06.6X2A	Traumatic subarachnoid hemorrhage with loss of consciousness of 31 minutes to 59 minutes, initial encounter	ICD-10-CM
S06.6X3A	Traumatic subarachnoid hemorrhage with loss of consciousness of 1 hour to 5 hours 59 minutes, initial encounter	ICD-10-CM
S06.6X4A	Traumatic subarachnoid hemorrhage with loss of consciousness of 6 hours to 24 hours, initial encounter	ICD-10-CM
S06.6X5A	Traumatic subarachnoid hemorrhage with loss of consciousness greater than 24 hours with return to pre-existing conscious level, initial encounter	ICD-10-CM
S06.6X6A	Traumatic subarachnoid hemorrhage with loss of consciousness greater than 24 hours without return to pre-existing conscious level with patient surviving, initial encounter	ICD-10-CM
S06.6X7A	Traumatic subarachnoid hemorrhage with loss of consciousness of any duration with death due to brain injury prior to regaining consciousness, initial encounter	ICD-10-CM
S06.6X8A	Traumatic subarachnoid hemorrhage with loss of consciousness of any duration with death due to other cause prior to regaining consciousness, initial encounter	ICD-10-CM
S06.6X9A	Traumatic subarachnoid hemorrhage with loss of consciousness of unspecified duration, initial encounter	ICD-10-CM



Appendix C. Specifications for Parameters for this Request

Sentinel's Cohort Identification and Descriptive Analysis (CIDA) tool version 5.0.5 was used to compare the frequency of diagnoses for intracranial hemorrhage using International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM) and International Classification of Diseases, Tenth Revision, Clinical Modification (ICD-10-CM) codes.

Enrollment Gap: 45 days

Age Groups: 21-64, 65-74, 75-84, 85-99 years

Enrollment Requirement: 183 days for incidence scenarios; 0 days for prevalence scenarios

Coverage Requirement: Medical and drug coverage

Event

Scenario	Query Start Date	Query End Date	Event	Event Code Type	Incident with Respect To:	Washout (days)	Cohort Definition	Care Setting
1	4/1/2015	9/30/2015	Intracranial Hemorrhage	ICD-9-CM	ICD-9-CM	183	First valid event only	Any
2	4/1/2016	9/30/2016	Intracranial Hemorrhage	ICD-10-CM	ICD-10-CM	183	First valid event only	Any
3	10/1/2010	9/30/2016	Intracranial Hemorrhage	ICD-9-CM	ICD-9-CM or ICD-10-CM	183	Any valid events	Any
4	10/1/2010	9/30/2016	Intracranial Hemorrhage	ICD-10-CM	ICD-9-CM or ICD-10-CM	183	Any valid events	Any
5	10/1/2010	9/30/2016	Intracranial Hemorrhage	ICD-9-CM	N/A	0	Any valid events	Any
6	10/1/2010	9/30/2016	Intracranial Hemorrhage	ICD-10-CM	N/A	0	Any valid events	Any

ICD-9-CM and ICD-10-CM are provided by Optum360. ICD-10-CM codes were mapped from ICD-9-CM codes using the Centers for Medicare and Medicaid Services General Equivalence Mappings.